


\$2.50

47

Popular Computing

February 1977 Volume 5 Number 2



Sixteen points are located on a 100 x 100 grid, as shown on the cover. The points are to move as follows:

Each point moves $\frac{1}{7}$ of the way toward the next higher numbered point. Thus, for example, point No. 5 should move to a new position given by:

$$x = 58 + \frac{1}{7}(75 - 58)$$


$$y = 22 + \frac{1}{7}(51 - 22)$$

which puts it at (60.42857, 26.14286). Point No. 16 moves in the direction of point No. 1. The set of points is to be moved in order, from No. 1 through No. 16, with the process then repeating, starting again with point No. 1.

It seems intuitively clear that the process will converge; that is, all the points will eventually meet.

If so, where?

Will it be close to the center of the initial set of 16 points; that is, at the average of the initial x and y values, which is 55.3125 and 48.6875?



POPULAR COMPUTING is published monthly at Box 272, Calabasas, California 91302. Subscription rate in the United States is \$20.50 per year, or \$17.50 if remittance accompanies the order. For Canada and Mexico, add \$1.50 per year to the above rates. For all other countries, add \$3.50 per year to the above rates. Back issues \$2.50 each. Copyright 1977 by POPULAR COMPUTING.

Editor: Audrey Gruenberger
 Publisher: Fred Gruenberger
 Associate Editors: David Babcock
 Irwin Greenwald

Contributing editors: Richard Andree
 William C. McGee
 Thomas R. Parkin

Advertising Manager: Ken W. Sims
 Art Director: John G. Scott
 Business Manager: Ben Moore

Reproduction by any means is prohibited by law and is unfair to other subscribers.

Knuth 2

An Approach to Floyd's Problem

Last month I discussed some of the virtues of Floyd's problem, which is to divide the numbers

$$\sqrt{1}, \sqrt{2}, \dots, \sqrt{50}$$

into two parts whose sums are as equal as you can make them after ten seconds of computer time. Here is the way I decided to approach the subject; perhaps some reader will have a much better idea.

Since $\frac{1}{2}(\sqrt{1} + \sqrt{2} + \dots + \sqrt{50})$ is equal to

119.51790 03017 60392 24702 02231... ,

we seek the subset of $\{\sqrt{1}, \sqrt{2}, \dots, \sqrt{50}\}$ whose sum least exceeds this value.

In the first place it is helpful to estimate the kind of results that might be expected. Most subsets of the given set have about 25 elements, and in fact the number of subsets with exactly 25 elements is $50!/25!25!$; this is about

$$\frac{2^{50}}{\sqrt{25\pi}},$$

according to Stirling's approximation. [The exact number is 12641 06064 37752, compared to

$$2^{50} = 1\,12589\,99068\,42624.]$$

All of these subsets have a sum which lies between

$$\sqrt{1} + \sqrt{2} + \dots + \sqrt{25} \approx 85.6 \quad \text{and} \quad \sqrt{26} + \sqrt{27} + \dots + \sqrt{50} \approx 153.4,$$

and they tend to cluster about the average value 119.5. So we have more than 10^{14} numbers packed into an interval of length less than 70; therefore $70/10^{14}$ appears to be a conservative estimate for the amount by which the best partitioning exceeds

$$\frac{1}{2}(\sqrt{1} + \sqrt{2} + \dots + \sqrt{50}).$$

But not all of these subsets will give different sums. In the first place, the seven numbers

$$\sqrt{1}, \sqrt{4}, \dots, \sqrt{49} = 1, 2, \dots, 7$$

will yield only integer values (and it is easy to verify that each integer from 0 to 28 can be represented). Our problem therefore reduces to finding a subset of the 43 numbers

$$\{\sqrt{2}, \sqrt{3}, \sqrt{5}, \sqrt{6}, \dots, \sqrt{48}, \sqrt{50}\}$$

whose sum has a fraction part least exceeding

$$.51790\ 03017\ 60392\ 24702;$$

unless the integer part of the sum is unusually large or small, we will be able to adjust it by adding a suitable subset of $\{1, 2, \dots, 7\}$, bringing the integer part up to 119.

We now have 2^{43} possibilities to try; but they aren't all distinct, either. The values

$$\{\sqrt{2}, \sqrt{8}, \sqrt{18}, \sqrt{32}, \sqrt{50}\} = \{\sqrt{2}, 2\sqrt{2}, 3\sqrt{2}, 4\sqrt{2}, 5\sqrt{2}\}$$

lead to subsets whose sum is restricted to sixteen values

$$0, \sqrt{2}, 2\sqrt{2}, \dots, 15\sqrt{2},$$

so only 16 of the 32 subsets are different. Similarly, there are only eleven essentially different subsets of

$$\sqrt{3}, \sqrt{12}, \sqrt{27}, \sqrt{48}$$

to try. By means of these observations our program can do in 10 seconds what it would take $(10 \cdot 32 \cdot 16) / (16 \cdot 11) \approx 29$ seconds to do otherwise.

But there are still more than $2^{41} > 10^{12}$ possibilities remaining, and this is far too large; if it takes $100\mu\text{s}$ for me to test one subset, I will have time to test only 10^5 subsets.

One way to gain speed is to divide

$$\{\sqrt{2}, \sqrt{3}, \dots, \sqrt{48}, \sqrt{50}\}$$

into two parts A and B, and to form a table containing all sums of the subsets of A. Then for each subset of B we can look in the table to see if there is an entry with the appropriate leading bits.

That is what I eventually did. Let

$$A = \{\sqrt{2}, \sqrt{8}, \sqrt{18}, \sqrt{32}, \sqrt{50}, \sqrt{3}, \sqrt{12}, \sqrt{27}, \\ \sqrt{48}, \sqrt{5}, \sqrt{6}, \sqrt{7}, \sqrt{10}, \sqrt{11}, \sqrt{13}, \sqrt{14}, \sqrt{15}\},$$

so that the subsets of A take on $(11/16) \cdot 2^{16}$ different values. I stored the 32 leading bits of these fraction parts into a hash table of size 2^{16} , and in another table of size 2^{16} I stored a bit pattern identifying the corresponding subset. (The hash code was the low order 16 bits of each 32 bit key.) Let $B = \{\sqrt{17}, \sqrt{19}, \sqrt{20}, \dots, \sqrt{47}\}$ be the remaining set of 26 elements; in the machine I used only the 32 leading bits of the fraction parts of each number \sqrt{k} . Since my hash table contained $(11/16) \cdot 2^{16}$ more or less random 32-bit numbers, I expected to get a match once out of every $2^{32}/((11/16) \cdot 2^{16}) = (16/11) \cdot 2^{16}$ times I looked up another 32-bit value. Therefore I wanted to have a program that generated 2^{17} or so subsets S of B, looking up the 32 bits corresponding to $(.51790 \dots - \sum S) \bmod 1$ in the hash table. I figured that would probably give me two or three matches, and I could choose the best corresponding partition. But I would have only about 75 to 80 μ s to spend per subset, so I needed a fast way to probe the hash table and to generate the subset sums.

The solution was to use ordered hash tables with linear probing; this is a variant of hashing which Ole Amble and I had discovered a few years ago [1]. Such hash tables are especially well suited to cases when searches are unsuccessful, requiring only 2.1 probes per search in this case. For the subset generation, I used Gray code [2], since this meant that only one subset element changed state each time and the subset sum was therefore easy to update. It took about 1.5 seconds to build the hash table. I started the subset generation of B in a random part of its cycle, since I knew that I would only be able to look at a small fraction of its subsets.

The best of the three results I got before 10 seconds expired was actually typed out after only 6 seconds, namely

$$\begin{aligned} & \sqrt{2} + \sqrt{3} + \sqrt{4} + \sqrt{5} + \sqrt{6} + \sqrt{12} + \sqrt{17} + \sqrt{19} + \sqrt{22} + \sqrt{27} + \\ & \sqrt{28} + \sqrt{29} + \sqrt{33} + \sqrt{34} + \sqrt{35} + \sqrt{37} + \sqrt{38} + \sqrt{49} + \sqrt{50} \\ & \approx 119.51790 \ 03021 \ 65123 \ 39726 \ 54768 \ . \end{aligned}$$

This sum differs from its complementary sum by approximately $8 \cdot 10^{-10}$. (The optimum partition might well be a thousand times better, as mentioned above; my program could have found it if 10 thousand seconds were available instead of ten!)

I used a fairly large and fast computer, the DEC KL-10 at Stanford's Artificial Intelligence Laboratory. If I had used a slower machine, I would have cut down the number of bits in each hash table entry from 32 to something less, one bit per factor of two in speed. If I had used a smaller machine, I would have had to make the hash table smaller.

If I really wanted to get the optimum partition, I think it could be found in about 30 minutes. The best approach I can think of would be to divide the 43 irrational numbers into two subsets having respectively $11/16 \cdot 2^{21}$ and 2^{21} distinct sums. I would sort each of these files of sums by their fraction parts; since each file contains about two million entries, this would be the time-consuming part of the operation. (Instead of using a general-purpose sort routine I would write a special one, since it is easy to go from the sorted file for set A to the twice-as-long sorted file for $A \cup \{a\}$ by essentially merging the first file with itself. The total time to build the sorted set for A when A has n elements is therefore of order 2^n , while a general-purpose sort routine would take order $n \cdot 2^n$ steps.) Finally, given two large sorted files

$$x_1 \leq \dots \leq x_n \quad \text{and} \quad y_1 \leq \dots \leq y_n$$

and a number z, there is a nice algorithm which computes

$$\min\{x_i + y_j \mid x_i + y_j \geq z\}.$$

The reader will enjoy discovering this algorithm for himself; curiously it is essentially the same as Hamming's "p/q" algorithm in PC41-4, under a logarithmic transformation, with $x_p = \log p$ and $y_q = \log(qx)$.

I tested my hashing program by trying it first on

$$\{\sqrt{1}, \sqrt{2}, \dots, \sqrt{30}\}$$

and then on

$$\{\sqrt{1}, \sqrt{2}, \dots, \sqrt{40}\}.$$

I gave the optimum partition for the former case in the previous article (see issue No. 46); and I think I found the optimum partition also in the latter case:

$$\begin{aligned} &\sqrt{2} + \sqrt{5} + \sqrt{7} + \sqrt{13} + \sqrt{14} + \sqrt{17} + \sqrt{18} + \sqrt{19} + \sqrt{22} + \sqrt{23} + \\ &\sqrt{24} + \sqrt{26} + \sqrt{27} + \sqrt{29} + \sqrt{30} + \sqrt{32} + \sqrt{34} + \sqrt{38} + \sqrt{39} \\ &\qquad \qquad \qquad \approx 85.807894002346, \end{aligned}$$

$$\sqrt{1} + \sqrt{3} + \sqrt{4} + \sqrt{6} + \sqrt{8} + \sqrt{9} + \dots + \sqrt{40} \approx 85.807894002154.$$

However, I am not absolutely sure that this is the best, because there is a slight chance that an unusual rounding error might have occurred somewhere in the calculations.

I still think that the 10-second-limited problem is more interesting; according to Hamming's famous aphorism, the purpose of computing is insight, and I believe this problem brings some valuable insights into view.

Postscript: R. L. Graham recently told me about the following sets of nine square roots whose sums agree to 54 decimal places:

$$\begin{aligned} &\sqrt{100000001} + \sqrt{100000025} + \sqrt{100000031} + \sqrt{100000084} + \\ &\sqrt{100000087} + \sqrt{100000134} + \sqrt{100000158} + \sqrt{100000182} + \\ &\sqrt{100000198} \end{aligned}$$

$$\begin{aligned} &= 9000.00449 \ 99983 \ 56751 \ 33987 \ 35593 \ 13014 \ 12703 \ 30519 \\ &\quad 82221 \ 56985 \ 62024 \ 86633 \ \dots \end{aligned}$$

$$\begin{aligned} &\sqrt{100000002} + \sqrt{100000018} + \sqrt{100000042} + \sqrt{100000066} + \\ &\sqrt{100000113} + \sqrt{100000116} + \sqrt{100000169} + \sqrt{100000175} + \\ &\sqrt{100000199} \end{aligned}$$

$$\begin{aligned} &= 9000.00449 \ 99983 \ 56751 \ 33987 \ 35593 \ 13014 \ 12703 \ 30519 \\ &\quad 82221 \ 56985 \ 62028 \ 18260 \ \dots \end{aligned}$$

To find the secret of his construction, see Hardy and Wright's Theory of Numbers, 4th edition, p. 338 (notes on Section 21.10).

References

- [1] Ole Amble and Donald E. Knuth, "Ordered hash tables," The Computer Journal 17 (May, 1974), 135-142.
- [2] Martin Gardner, "Mathematical Games," Scientific American 227 (August, 1972), 106-109.



2-3-5 again

In issue No. 10 there first appeared this problem:

Show the logic of generating, in order, all numbers having only the factors 2, 3, and/or 5. The first 30 such numbers are:

2, 3, 4, 5, 6, 8, 9, 10, 12, 15, 16,
18, 20, 24, 25, 27, 30, 32, 36, 40,
45, 48, 50, 54, 60, 64, 72, 75, 80, 81.

The problem was attributed to Richard Hamming.

In issue No. 16, three different solutions to the 2-3-5 problem were shown, including one by Hamming. The following comes from Norman Sanders, Trondheim, Norway.

I had the problem of generating thousands of numbers of the form



$$2^a \cdot 3^b, \quad 2^a \cdot 3^b \cdot 5^c, \quad 2^a \cdot 3^b \cdot 5^c \cdot 7^d$$

in connection with the numerical validation of an approximation in prime-number theory. But that was in the mid-50's on EDSAC-1, which had a 1K store, no tapes, no divide, 1.5 millisecond cycle time, and zero-generation reliability. A divide subroutine was a time-consuming luxury that ruled out the first two solutions shown in PC16. The severe storage limitations ruled out long lists, and the low availability of the computer implied a simple program.

As an example of the general method, the solution for

$$2^a \cdot 3^b \cdot 5^c$$

was to generate three lists, A, B, and C (of which C consisted of a single entry only) and to transfer elements from list to list, freeing space at each transfer and each print instruction. In this way, I was able to exceed 2^{100} without reaching the limitation of the store.



Norman Sanders' algorithm
for the 2-3-5 problem.

Snapshot at the time of printing N = 25:

<u>PRINT LIST</u>	<u>A-LIST</u>	<u>B-LIST</u>	<u>C-LIST</u>
2	1 0 0	0 1 0	0 0 1
3	0 1 0	0 0 1	0 0 2
4	2 0 0	0 2 0	0 0 3
5	0 0 1	0 1 1	0 0 4
6	1 1 0	0 0 2	
8	3 0 0	0 3 0	
9	0 2 0	0 2 1	
10	1 0 1	0 1 2	
12	2 1 0	0 4 0	
15	0 1 1	0 0 3	
16	4 0 0	0 3 1	
18	1 2 0		
20	2 0 1		
24	3 1 0		
25	0 0 2		
	0 3 0		
	1 1 1		
	5 0 0		
	2 2 0		
	3 0 1		
	0 2 1		
	4 1 0		

The A-list consists of triples (a, b, c) , whose values are $2^a \cdot 3^b \cdot 5^c$, in increasing order of value. The B-list consists of triples $(0, \beta, \delta)$, whose values are $3^\beta \cdot 5^\delta$, in order. The C-list contains the single triple $(0, 0, \Gamma)$, of value 5^Γ . Initially the A-list consists of the single entry $(1, 0, 0)$; the B-list of the single entry $(0, 1, 0)$, and C of the entry $(0, 0, 1)$. The logic of the algorithm is shown in the flowchart.

The entries in the lists above the dotted lines have been removed. $N = 25(0, 0, 2)$ has just been printed, and the next step is to compare $(1, 0, 2)$ ($= 50$) with $(0, 1, 2)$ ($= 75$) at the top of the B-list. It is less, so $(1, 0, 2)$ will be appended below $(4, 1, 0)$ in the A-list.

A few pertinent remarks:

Don't throw anything away. I'm far too experienced today to reproduce such a simple method.

Document your stuff so that you can understand it twenty years later. You never know what the Hamming birds are going to come up with.

When we had hardly any computers, the algorithms we used had to treat the idiosyncracies of the hardware very carefully. Now that we have thousands of computers to choose from, most programming seems to be machine-independent. Computing has become Aristotelian.

We managed to get a lot of useful work out of the primitive computers. If we hadn't they would have stayed primitive, or died out all together.

[Mr. Sanders is the author of The Corporate Computer; How to Live With an Ecological Intrusion, McGraw-Hill, 1973. His pixie-ish lack of respect for authority should not detract from his computing wisdom.]

In Martin Gardner's "Mathematical Games" column in the November 1976 Scientific American, he shows that the following trick works:

Ask someone to pick a number, write it down, and show it to you. You add one to his number and divide by some other number that you have chosen (and secretly written down) and give him the result. Your friend now has two numbers and is to follow this plan: Given two numbers, add one to the second number, divide by the first number, and record the result as the third number. Then, using the last two of these, repeat the procedure two more times; the final result will be your hidden number.

The arithmetic for the choice of 17 by the victim and 1.2345679 as your hidden number is shown here:

- ① Your hidden number: 1.2345679
- ② His first number: 17
- ③ You give him $\frac{18}{1.2345679} = 14.58$
- ④ $\frac{15.58}{17} = .9164705882$
- ⑤ $\frac{1.9164705882}{14.58} = .1314451707$
- ⑥ $\frac{1.1314451707}{.9164705882} = 1.2345679$

It is easy to show that this must work. The successive stages for any two starting numbers will be:

- | | |
|-------------------|----------------------|
| ① x | ② y |
| ③ $\frac{y+1}{x}$ | ④ $\frac{y+x+1}{xy}$ |
| ⑤ $\frac{x+1}{y}$ | ⑥ x |

Thus, the numbers will repeat on a cycle of 5.

Problem: To find a simple similar algorithm that will repeat on a cycle of 6.



Problem Solution


PC47-13

Wendy's Problem (Number 148, PC45-6) called for counting all possible triangles that can be formed on a 10 x 10 grid of points and tabulate the results by areas.

Associate Editor David Babcock produced the tabulation shown here. The column labelled N is twice the area and column K shows how many such triangles there are. The total of the K values is 161,700.

It is of interest to note that there are 4448 combinations of 3 points that do not form a triangle, and the curious distribution of the other 80 possibilities.

N	K	N	K	N	K	N	K	N	K
0	4448	1	5852	2	8376	3	7236	4	9064
5	5816	6	9464	7	4940	8	7812	9	5404
10	6128	11	3160	12	7904	13	2828	14	4760
15	4436	16	5020	17	2260	18	5032	19	2016
20	4160	21	3136	22	2416	23	1600	24	4368
25	1924	26	1888	27	1916	28	2640	29	1112
30	2640	31	976	32	1972	33	1116	34	1120
35	1420	36	1960	37	664	38	840	39	736
40	1300	41	504	42	1336	43	432	44	624
45	692	46	448	47	320	48	932	49	500
50	320	51	296	52	304	53	184	54	480
55	152	56	524	57	176	58	144	59	104
60	168	61	88	62	88	63	260	64	192
65	44	66	72	67	40	68	48	69	48
70	32	71	24	72	148	73	16	74	8
75	16	76	8	77	12	78	8	79	8
80	4	81	36						



F-N SEQUENCES

In the sequences shown in Table A, F-1 is the familiar Fibonacci sequence, in which

$$F_n = F_{n-1} + F_{n-2}$$

It is well known that the ratio of one term to the preceding term

$$\text{converges to } \frac{F_k/F_{k-1}}{2} = \frac{1 + \sqrt{5}}{2} = 1.618033988\dots$$

The sequence F-2, starting with three ones, is formed by the relation

$$F_n = F_{n-2} + F_{n-3}$$

and it appears to converge to the ratio 1.3247... We say "appears to converge" because the ratio of one term to the preceding term has a cycle of two values. For example, successive ratios for the first few terms are:


16/12	=	1.3333	}
21/16	=	1.3125	
28/21	=	1.3333	}
37/28	=	1.3214	
49/37	=	1.3243	}
65/49	=	1.3265	
86/65	=	1.3231	}
114/86	=	1.3256	
151/114	=	1.3246	}
200/151	=	1.3245	

so that the value to which the ratio converges is the average of two ratios. Similarly, for F-3, the convergent value will be found by averaging three successive ratios.

F-1	F-2	F-3	F-4	F-5
1	1	1	1	1
1	1	1	1	1
2	1	1	1	1
3	2	1	1	1
5	2	2	1	1
8	3	2	2	1
13	4	2	2	2
21	5	3	2	2
34	7	4	2	2
55	9	4	3	2
89	12	5	4	2
144	16	7	4	3
233	21	8	4	4
377	28	9	5	4
610	37	12	7	4
987	49	15	8	4

Table A

The first 16
terms of the
first five
F-N sequences.



Subsequent members of the F-N family of sequences are defined similarly:

$$F-3: F_n = F_{n-3} + F_{n-4} \quad \text{converging to } 1.2207\dots$$

$$F-4: F_n = F_{n-4} + F_{n-5} \quad \text{converging to } 1.1672\dots$$

$$F-5: F_n = F_{n-5} + F_{n-6} \quad \text{converging to } 1.1347\dots$$

$$F-6: F_n = F_{n-6} + F_{n-7} \quad \text{converging to } 1.1128\dots$$

For the sequences F-1 through F-8, the convergent values plot as shown. What is the curve that best expresses the relationship shown in the curve? Is the curve asymptotic to $y = 1$?

PROBLEM 161



47
N-SERIES

Log 47	1.672097857935717464414219399449200640159803098429948
ln 47	3.850147601710058586820950669772173708896050502020224
$\sqrt{47}$	6.855654600401044124935871449084848960460643461001326
$\sqrt[3]{47}$	3.608826080138694689252517295958892614905551690162338
$\sqrt[10]{47}$	1.469636013359929253623112224371627488588820827562794
$\sqrt[100]{47}$	1.039252262310962953384796629222198650121736878483157
e^{47}	258131288619006739623.2858002152733804316370829930440 6081061397243682275040978221987
π^{47}	232297222236041657886385.7994890281440584469844961463 1767505405499833821420635585
$\tan^{-1} 47$	1.549522940770835415559197706584603144431776847570469

Cunningham's Process

In the May, 1962 American Mathematical MONTHLY, G. S. Cunningham showed that any positive fraction can be represented as a sum of distinct unit fractions.

$$\frac{a}{b} = \frac{1}{b} + \underbrace{\frac{1}{b} + \frac{1}{b} + \cdots + \frac{1}{b}}_{a \text{ terms}}$$

Retain one of the duplicating terms. Replace each of the others by the identity

$$\frac{1}{n} = \frac{1}{n+1} + \frac{1}{(n+1)n}$$

and repeat this process until all terms are distinct.

For example:

$$\begin{aligned} \frac{2}{3} &= \frac{1}{3} + \frac{1}{3} \\ &= \frac{1}{3} + \frac{1}{4} + \frac{1}{12} \end{aligned}$$

$$\begin{aligned} \frac{3}{4} &= \frac{1}{4} + \frac{1}{4} + \frac{1}{4} \\ &= \frac{1}{4} + \frac{1}{5} + \frac{1}{20} + \frac{1}{5} + \frac{1}{20} \\ &= \frac{1}{4} + \frac{1}{5} + \frac{1}{20} + \frac{1}{6} + \frac{1}{30} + \frac{1}{21} + \frac{1}{420} \end{aligned}$$

$$\begin{aligned}
\frac{4}{5} &= \frac{1}{5} + \frac{1}{5} + \frac{1}{5} + \frac{1}{5} \\
&= \frac{1}{5} + \frac{1}{6} + \frac{1}{30} + \frac{1}{6} + \frac{1}{30} + \frac{1}{6} + \frac{1}{30} \\
&= \frac{1}{5} + \frac{1}{6} + \frac{1}{30} + \frac{1}{7} + \frac{1}{42} + \frac{1}{31} + \frac{1}{930} + \frac{1}{7} + \frac{1}{42} + \frac{1}{31} + \frac{1}{930} \\
&= \frac{1}{5} + \frac{1}{6} + \frac{1}{30} + \frac{1}{7} + \frac{1}{42} + \frac{1}{31} + \frac{1}{930} + \frac{1}{8} + \frac{1}{56} + \frac{1}{43} + \frac{1}{43 \cdot 42} \\
&\quad + \frac{1}{32} + \frac{1}{32 \cdot 31} + \frac{1}{931} + \frac{1}{931 \cdot 930} \\
&= \frac{1}{5} + \frac{1}{6} + \frac{1}{30} + \frac{1}{7} + \frac{1}{42} + \frac{1}{31} + \frac{1}{930} + \frac{1}{8} + \frac{1}{56} + \frac{1}{43} + \frac{1}{1806} \\
&\quad + \frac{1}{32} + \frac{1}{992} + \frac{1}{931} + \frac{1}{865830}
\end{aligned}$$

In each example, the fraction used was $\frac{N-1}{N}$

Problem: extend this table:

N	Number of final terms	Largest denominator
3	3	12
4	7	420
5	15	865830



ANNUAL INDEX

Volume 4 1976
Issues 34 through 45
230 pages

PC47-19

2-5 problem 45-8
3x+1 problem 45-14
18 Symposia 44-4
A to the B problem 42-11
Algebra Guidelines 44-17
And Tomorrow..The World 43-6
Andree, Prof. Richard 45-8
Another Way 36-18
Armer, Paul 44-6, 45-15
Armerding, George 42-17
Art of Computing essays:
 Bracketing 35-5
 Using New Tools 40-4
 Assemblers 41-8
APF Mark 51 34-16
Atanasoff, John 43-9
Assemblers 41-8
Assemblers II 42-9
Automated Mastermind 44-7
Babcock, David 38-5, 39-3, 42-2, 43-19
Backtracking problem 34-1
Backtracking II article 34-3, 39-17
Bank check problem 37-7
Beeler, Mike 36-10, 41-15
Beltran, Sergio 44-5
Berger, Robert 45-15
Bennett, William 43-17
Bicentennial star problem 36-1
Blankenbaker, John 39-14
Bracketing 35-5, 39-13, 44-8
Braiding problem 45-11
Breuer, Shlomo 36-15
Cady, Dorothy 34-10
Cald's Sequence 41-16
Centafactorials 39-20
Circle partitioning problem 36-8
Circuitous race problem 38-18
Circulating pump problem 45--5
Coleman, C. B., Jr. 44-3
Combinations subroutine contest 38-9
Compound interest 34-13
Computational Mathematics 36-15
Computer Applications 43-17
Computer Chess 39-7
Computer Industry Association 43-6
Concerning File Management Systems 43-1
Constrained random walk problem 42-1
Contest 4 35-2
Contest 5 36-4
Contest 2 outcome 36-10
Contest 6 37-4
Contest 7 38-6
Contest 3 outcome 38-9

Contest 8 39-5
Contest 9 40-3
Contest 4 outcome 40-12
Contest 10 41-2
Contest 5 outcome 41-14
Contest 11 42-4
Contest 6 outcome 42-8
Contest 8 outcome 44-20
Contest 12 43-10
Contest 7 outcome 43-14
Contest 13 44-1
Contest 9 outcome 44-13
Contest 14 45-1
Countdown problem 43-10
Counting string lengths 34-18, 35-17
Creative Computing 34-9
Croy, Timothy 44-18
CSR function 34-10, 35-3
Cube route solution 38-5
Cubes with primes 40-11
Daughterity, Walter 36-10
Dialogue 41-3
Duff, Tom 43-14, 44-18
Eckert, J. Presper 44-4
Eight dice problem 39-17
Error amplification problem 40-6
Factorials cycles 36-3
Faulty flowchart 44-9
Ferguson, David 37-4
Ferris, John 38-9
Fibonacci table 34-14
File Management systems 43-1
Forecast I 37-1
Four-square ratio problem 45-10
Foy, Nancy 43-6
Games With the Pocket Calculator 39-10
Gammadion 41-1
gcd problem 45-4
Gemero problem 40-19
Generating triangles problem 45-1
Gilbreath conjecture 42-17
Giving a Speech on Computing 43-11
Goldstine, Herman 43-9
Gordon, Barry 45-15
Graham, Neill 44-19
Greenwald, Irwin 42-9, 45-15
Grosch, H. R. J. 43-7, 44-4
Gruenberger, Fred 41-3, 38-5, 42-17, 43-1, 45-15
Guessword puzzles 35-14
Guidelines for algebra 44-17
Halt button on SR-52 40-14
Hamming, Richard 36-16, 40-5, 41-3, 44-4

Hawkes, Pamela 36-17
 Herschfeld, Aaron 35-3
 HP-65 Users club 36-11
 Hurley, James 42-11
 IBM card problem solution 37-16
 K-Column Fibonacci problem 40-1, 44-13
 K-Level Sieve problem 38-6, 43-14, 44-18
 K-Moduli Sequence problem 44-1
 Kallin, Sten 39-19
 Katzan, Harry, Jr. 35-5
 KENBAK-1 39-14
 Killgrove, R. B. 42-17
 Knuth, Donald E. 38-8
 Koetke, Walter 37-4
 Lander, L. J. 42-17
 Life or Death problem 37-14
 Malik, Rex 43-6
 Mastermind 35-14
 McCracken, Daniel 45-15
 Merging problem solution 34-16
 Mersel, Jules 37-8
 The Mind Tool 44-19
 Minimum Fortran problem 42-4
 Motil, John 39-6
 N in ones and zeros 37-4, 42-8
 Nelson, Richard 36-11
 New irrational number 39-3
 Newborn, Monroe 39-7
 Nutt, Roy 41-8
 One key calculator problem 45-9
 Ones and zeros make N 37-4, 42-8
 Outguess contest 39-4
 Overdue--or Overdone? 41-18
 Parkin, Thomas R. 34-3, 36-5, 37-16,
 42-17, 44-13
 Periodicals list 45-2
 Peripatetic Jumping Bean 38-1
 Pocket calculator problems 38-16, 45-9
 Poland, Clarence 45-15
 Prime Path problem 45-7
 Primes Sieve problem 36-4
 Problem definition 38-14
 Programming Class Project 45-11
 Proprietary Software 44-3
 Q problem solution 36-5
 Quick and Dirty RNG 38-8
 Quick, J. H. 36-12
 Ralston, K. E. 42-17
 Random Number Generator 43-19
 Random numbers by division 38-8
 Raphael, Bertram 44-8
 Recurse program 39-6

Redelmeier, Hugh 43-14, 44-18
 Ring-a-Ding problem 39-1
 Robinson, Herman P. 34-11, 35-3
 Sandfelder, M. E. 40-18
Scientific Analysis on the Pocket
 Calculator 39-9
 Schwartz, Mordecai 42-5, 42-12, 43-19
 Schwartz's random number generator 43-19
 Sense switch on SR-52 40-15
 Sequences 42-5
 Shallit, Jeffrey 40-12
 Shortcut problem 36-14
 Simpson's Rule 40-6
 Smith, Jon M. 39-9
 Smith, Richard D. 38-15
 Slowgrow correction 37-12
 Snoopy's Water Dish problem 35-12
 Square Spiral problem 35-1, 40-12
 SR-50 rating 34-16
 SR-52 Notes 40-14, 41-20
 SR-52 Review 38-3
 SR-52 Sense Switch 40-15
 SR-52 Software 44-3
 Stack Action problem 37-3, 40-18
 Stolovitch, Harold 39-10
 Storage Filling problem 34-12
 Sum Fun Puzzle Club 44-7
 Sum of One problem 36-18
 Summing 7-Card Decks problem 40-16
 Swiertz, Donald 44-20
 Symposium 18 transcript 45-15
 Take/Skip problem 38-6
 Tanaka, Richard 45-15
 Thiagarajan, Sivasailam 39-10
 The Thinking Computer 44-8
 Thompson, Harold 39-20
 Three X Plus One 45-14
 Turbo-Encabulator 36-12
 Up Your Word game 37-13
 Using New Tools article 40-4
 von Neumann, John 43-9
 Wagner, Sam 42-8
 Weinberger, Jared 37-12
 Weizenbaum, Joseph 43-18, 45-15
 Wendy's problem 45-6
 White, Robert 45-15
 Why Are You Waiting? poster 36-17
 Winslow, Leon 40-5
 X to the X to the X 39-13
 Yardley, Herbert O. 43-18
 Z-Sequence 42-12
 Zuse, Konrad 43-9
 Zwas, Gideon 36-15